

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁵ : C12N 9/00, 9/42	A1	(11) International Publication Number: WO 91/18974 (43) International Publication Date: 12 December 1991 (12.12.91)
(21) International Application Number: PCT/US91/03756 (22) International Filing Date: 29 May 1991 (29.05.91) (30) Priority data: 529,655 29 May 1990 (29.05.90) US (60) Parent Application or Grant (63) Related by Continuation US 529,655 (CIP) Filed on 29 May 1990 (29.05.90) (71) Applicant (for all designated States except US): CHEMGEN CORPORATION [US/US]; 16016 Industrial Drive, Gaithersburg, MD 20877 (US).		(72) Inventors; and (75) Inventors/Applicants (for US only) : FODGE, Douglas, Walter [US/US]; 7308 Righters Mill Road, Rockville, MD 20855 (US). ANDERSON, David, Martin [US/US]; 13509 Bailey Drive, Rockville, MD 20850 (US). PETTEY, Thomas, Michael [US/US]; 1400 Washingtonian Tower, 9701 Fields Road, Gaithersburg, MD 20878 (US). (74) Agents: BENT, Stephen, A. et al.; Foley & Lardner, 1800 Diagonal Road, Alexandria, VA 22313 (US). (81) Designated States: AT (European patent), AU, BE (European patent), BG, BR, CA, CH (European patent), DE (European patent), DK (European patent), ES (European patent), FI, FR (European patent), GB (European patent), GR (European patent), HU, IT (European patent), JP, KP, KR, LU (European patent), NL (European patent), NO, PL, RO, SE (European patent), US. Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
(54) Title: HEMICELLULASE ACTIVE AT EXTREMES OF pH AND TEMPERATURE AND THE MEANS FOR THE PRODUCTION THEREOF (57) Abstract Soil bacteria can be isolated which produce an enzyme capable of catalyzing the degradation of mannan-containing hemicellulose under conditions combining high pH and high temperature. Such bacteria can be cultured or used as sources of genetic information with which to engineer other microorganisms to produce the enzyme. Commercially useful quantities of native or recombinant hemicellulase can thus be produced by cultures consisting essentially of microorganisms capable of producing the enzyme.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	ES	Spain	MG	Madagascar
AU	Australia	FI	Finland	ML	Mali
BB	Barbados	FR	France	MN	Mongolia
BE	Belgium	GA	Gabon	MR	Mauritania
BF	Burkina Faso	GB	United Kingdom	MW	Malawi
BG	Bulgaria	GN	Guinea	NL	Netherlands
BJ	Benin	GR	Greece	NO	Norway
BR	Brazil	HU	Hungary	PL	Poland
CA	Canada	IT	Italy	RO	Romania
CF	Central African Republic	JP	Japan	SD	Sudan
CG	Congo	KP	Democratic People's Republic of Korea	SE	Sweden
CH	Switzerland	KR	Republic of Korea	SN	Senegal
CI	Côte d'Ivoire	LI	Liechtenstein	SU	Soviet Union
CM	Cameroon	LK	Sri Lanka	TD	Chad
CS	Czechoslovakia	LU	Luxembourg	TG	Togo
DE	Germany	MC	Monaco	US	United States of America
DK	Denmark				

- 1 -

HEMICELLULASE ACTIVE AT EXTREMES OF pH AND
TEMPERATURE AND THE MEANS FOR THE PRODUCTION THEREOF

Background of the Invention

Different enzymes are categorized as a
5 specific type of hemicellulase -- a glucanase, a
xylanase or a mannanase, for example -- based on an
ability to catalyze the hydrolysis of heteropoly-
saccharides composed of glucan, xylan or mannan,
respectively. It is known that enzymes that effect
10 hydrolysis of mannans, such as a galactan or a
glucomannan, are produced by various microorganisms,
including bacteria and fungi, and that they also
occur in some animals and in numerous plants. Among
the microorganisms that produce such mannanases are
15 species of *Aeromonas*, *Aspergillus*, *Streptomyces*,
Rhodococcus and *Bacillus*. See 160 METHODS IN
ENZYMOLGY Part A, Sect. II (1988).

Hemicellulases have been employed
commercially, for example, in the processing of
20 coffee, chocolate, cocoa, tea and cereals.
Processing these foodstuffs with hemicellulases
converts them to a more edible form, thereby
improving the energy value of the product, and
reduces solution viscosity. Hemicellulases have
25 thus been used to clarify fruit juices, to reduce
the viscosity of slurries or purees, to liquefy
certain cell wall solids and to modify taste.
Similarly, hemicellulases have also been employed to

- 2 -

mitigate problems associated with the presence of hemicellulose and other minor polysaccharides in pulps and in unprocessed cotton. By the same token, hemicellulose is a problem when it is present in sewage and industrial waste-water treatment, and in fracturing fluids used in the conditioning of oil-bearing well formations.

Some processes involved in commercial applications utilizing hemicellulose are characterized by conditions of high temperatures and high alkalinity. Hemicellulases in general, and mannanases heretofore described in particular, exhibit substantial activity at a pH range of 3-8 and temperatures below 60° C. These enzymes therefore cannot be used to process hemicellulose cost effectively in applications where these conditions are exceeded.

Nonspecific oxidizing chemicals have been employed to degrade hemicellulose in many commercial oil well fracturing solutions in the exceeded conditions. Incomplete degradation of the hemicellulose is effected, however, at all pHs and temperatures. This can prove to be inefficient, at best, thereby increasing processing costs and resulting in marginal usefulness. For example, partial degradation of mannan-based hydraulic gels, either by chemical oxidizers or by enzymes, leaves a residue that cannot be adsorbed into the surrounding formation. This may lead to a situation whereby the residue can coat fracture surfaces, thereby impeding oil exiting the formation. Accordingly, there is a need for a hemicellulase capable of retaining a substantial portion of its activity in conditions of high alkalinity and at temperatures exceeding 60° C. Such an enzyme would permit complete and efficient degradation of hemicellulose and modified hemicellulose under the

- 3 -

conditions that prevail in most of the commercial processes involving hemicellulose-based hydraulic gel.

Summary of the Invention

5 It is therefore an object of the present invention to provide a hemicellulase that displays substantial biological activity under conditions of high temperature and high alkalinity.

10 It is another object of the present invention to provide a method of isolating a microorganism capable of producing a heat and alkaline-resistant hemicellulase, as well as to provide a microbial culture consisting essentially of cells that produce the foregoing hemicellulase.

15 It is yet another object of the present invention to provide a method of producing commercially useful quantities of the foregoing hemicellulase.

20 A further object of the present invention is to provide a means for degrading mannan-based hemicelluloses under conditions of high alkalinity and elevated temperature such as are commonly found, for example, in the subterranean conditioning of oil fields, in the processing of some paper pulps and
25 certain food materials, and in sewage and industrial waste-water treatment.

30 In accomplishing the foregoing objects, there has been provided, in accordance with one aspect of the present invention, an enzyme in substantially pure form that catalyzes the degradation of hemicellulose and that has a pH profile for its hemicellulose-catalyzing activity that ranges from about pH 4.5 to about pH 11. In certain preferred embodiments, the pH profile of the enzyme peaks

- 4 -

between about pH 7 and about pH 9, and the enzyme catalyzes the degradation of a mannan-containing hemicellulose. In another preferred embodiment the enzyme is one that can catalyze the degradation of hemicellulose even when both pH and temperature are elevated into the range of 8-11 and to at least 60° C, respectively.

In accordance with another aspect of the invention, a microbial culture is provided that consists essentially of cells that produce an enzyme as described above. In a preferred embodiment, the microbial culture consists essentially of Gram-positive bacteria of a *Bacillus* species, such as *B. lentis*.

In accordance with other aspects of the present invention, an isolated polynucleotide is provided that codes for an enzyme that catalyzes the degradation of hemicellulose, said enzyme having a pH profile for its activity in catalyzing the degradation of hemicellulose that ranges from about pH 4.5 to about pH 11. Also provided are (i) a vector comprising a nucleotide sequence coding for such an enzyme and (ii) a host cell transformed with such a vector in order that the host cell expresses the enzyme.

In accordance with yet another aspect of the invention, a method of producing commercially useful quantities of the foregoing hemicellulase is provided, comprising the steps of:

(a) growing the microbial culture in a suitable broth medium;

(b) adding a mannan-containing hemicellulose to the broth culture during the growth phase or the stationary phase of the microorganism;

(c) obtaining a culture supernatant from the broth culture, the supernatant comprising a

- 5 -

predetermined concentration range of the hemicellulase;

(d) removing the hemicellulase by ultrafiltration;

5 (e) precipitating the hemicellulase with a precipitating agent selected from the group consisting of acetone and methanol; and

(f) recovering the hemicellulase.

10 In accordance with a further aspect of the invention, a method is provided for conditioning subterranean oil wells comprising the steps of:

(a) preparing a suspension comprised of water, a hydraulic gelling agent and a proppant, the proppant having an indicated acceptable mesh size;

15 (b) pumping an adequate amount of the suspension into an oil well under high pressure causing a fracture to occur in subterranean geologic formations adjacent to the well such that the suspension is deposited in the fracture;

20 (c) pumping an adequate amount of the foregoing hemicellulase with the suspension into the fracture to effect controlled degradation of the hydraulic gelling agent.

25 Other objects, features and advantages of the present invention will become apparent from the following detailed description. It should be understood, however, that the detailed description and the specific examples, while indicating preferred embodiments of the invention, are given by
30 way of illustration only since various changes and modifications within the spirit and scope of the invention will become apparent to those skilled in the art from this detailed description.

Brief Description of the Drawings

In drawings which illustrate embodiments of the invention,

Figure 1 presents representative data showing
5 the reduction in viscosity in relation to the amount
of time that enzyme dilutions from 14 liter
fermentations were incubated with the guar
solution.

Figure 2 is a graph which plots the activity
10 of a hemicellulase within the present invention
versus time (pH 9.0) at 60°, 75° and 90° C,
respectively.

Figure 3 shows the relative stability at
50° C over time of a dried lactose formulation of the
15 hemicellulase.

Figure 4 is a graph that compares the pH
profile of enzymatic activity for a hemicellulase of
the present invention versus a known, commercially
available enzyme.

20 Detailed Description of the Preferred Embodiments

It has been discovered that a hemicellulase
capable of catalyzing the degradation of mannan-
containing hemicellulose under conditions
characterized both by high alkalinity and by high
25 temperature can be produced using microorganisms
that synthesize the enzyme. Microorganisms
possessing the ability to produce a hemicellulase
within the present invention can be isolated by
conventional methods from the soil, where they
30 comprise a reproducible, finite subsection of
indigenous microflora. Hemicellulase-producing
microorganisms can also be produced by transforming
another microorganism of choice, such as *B. subtilis*

- 7 -

or *B. brevis*. Transformation may proceed with hemicellulase-encoding DNA obtained, via standard recombinant DNA techniques, from such indigenous soil microflora.

5 In cultures consisting essentially of microorganisms that produce a hemicellulase within the present invention, commercially significant quantities of the enzyme can be obtained by isolation techniques that are well known in the art.

10 In this description, a "culture consisting essentially of a specified type of microorganism," such as a hemicellulase-producing microorganism, is a culture that is comprised primarily of microbes of that type, to the extent that the salient functional

15 property or properties of the culture are determined by those microbes. But other types of microorganisms may be present, for example, in a culture consisting essentially of a hemicellulase-producing microorganism, so long as the other types

20 do not interfere significantly with hemicellulase production by the culture.

Microorganisms producing a hemicellulase with the foregoing properties can be isolated, pursuant to the present invention, from soil samples

25 collected from diverse geographical areas. The soil samples are collected primarily from the top two inches of soil and cultured in selective enrichment media.

Techniques for selectively isolating

30 microorganisms with certain desired characteristics are well known in the art. Microorganisms can be selected for the capacity to produce a particular enzyme by exposing samples potentially containing such microorganisms to conditions wherein

35 replication and growth are dependent upon the capacity to produce the desired enzyme. One widely used method of selection is exposing a sample

- 8 -

5 containing a large, diverse microbial population to
a medium consisting of a single carbon source. See
160 METHODS IN ENZYMOLOGY 180-86 (1988). Only those
microorganisms capable of producing enzymes that can
degrade the carbon source will be recovered by this
method. Selective culturing techniques of this sort
effectively separate microorganisms producing the
desired enzyme from the myriad of other
microorganisms comprising the normal flora of the
habitat sampled.

10 In a preferred embodiment of the present
invention, aliquots of soil samples are inoculated
into an alkaline culture medium, wherein
hemicellulose serves as the sole carbon source.
15 Dilutions of the sub-cultures grown in the same
medium are plated out onto a solid medium which also
contains hemicellulose as the sole carbon source.
Morphologically distinct colonies are isolated and
then screened for hemicellulase activity. In a
20 particularly preferred embodiment of this invention,
a selective enrichment broth at a pH range of 9-9.5,
wherein purified guar gum is the sole carbon source
is inoculated with a sample of soil, incubated at 37°
C and aerobically agitated.

25 After incubation, further dilutions of the
initial culture are made using the selective
enrichment broth. After several passages, a
dilution series of the most dilute broth culture is
made in normal saline and plated out on a solid
30 medium which contains guar as the sole carbon
source. After incubation at 34° C for 5-7 days,
morphologically distinct colonies are isolated from
the solid medium and screened for hemicellulase
activity.

35 The initial selection for microorganisms
which utilize hemicellulose as a sole carbon source
at an elevated pH results in the recovery of a

- 9 -

finite fraction of all soil microorganisms. Typically, about 40% of the isolates are characterized by an ability to produce a hemicellulase within the present invention. A particularly preferred subgroup of this fraction includes species of the Gram-positive genus *Bacillus*.

Exemplary of this preferred subgroup is a strain of *B. lentis*, designated CMG1240, the general properties of which are summarized in Table 1. Strain CMG1240 has been deposited, in accordance with the Budapest Treaty, at the American Type Culture Collection (Rockville, Maryland) under accession No. 55045. The key characteristic of strain CMG1240, as for any microbial strain of the preferred subgroup, is its ability to produce in culture medium a hemicellulase according to the present invention.

- 10 -

TABLE I

- a. Morphology
- 5 (1) bacillus shape; (0.8 X 1.7-2.3) μ m
 (2) occurs predominately in singles and pairs
 (3) capable of forming endospores
 (4) Gram-positive (also appears Gram-variable or Gram-negative)
- b. Growth Conditions
- 10 Tryptic soy agar plate culture: colonies are irregular, convex, smooth, undulate, butyrous, opaque and 2 mm in diameter (72 hours, 37° C).
- c. Biochemical Properties:
- 15 (1) Aerobic
 (2) Catalase positive
 (3) Oxidase negative
 (4) Non-motile
- 20 (5) Growth temperature: 30-50° C, maximum growth rate occurring at 41°C
 (6) Growth pH (cultured at 34° C at 350 rpm for 10 hours in a basal glucose-mineral salts medium): 7.0-8.5, maximum growth rate occurring at pH 7.5
- 25 (7) Casein and gelatin decomposition: negative
 (8) Starch hydrolysis: positive
 (9) Growth in 1-5% NaCl: positive
- 30 (10) Carbon sources utilized include: D-glucose, D-galactose, D-fructose, D-xylose, lactose, maltose, sucrose, alpha-cyclodextrin, dextrin, glycogen, N-acetyl-glucosamine, L-arabinose, cellobiose, gentiobiose, lactulose,
- 35 mannitol, D-mannose, D-melibiose, D-trehalose, turanose, A-ketobutyric acid, uridine, and m-inositol.
- (11) Hydrogen sulfide production: negative
 (12) Indole production: negative
- 40 (13) Utilization of citrate: negative
 (14) Urease: negative
 (15) Voges-Proskauer: negative
 (16) Phenylalanine deaminase: negative
- 45 (17) Lysine decarboxylase: negative
 (18) Ornithine decarboxylase: negative
 (19) Arginine dehydrolase: negative

Conventional methods for measuring cellulase/hemicellulase activities are described, for example, in 160 METHODS IN ENZYMOLOGY 180-86 &

50 368-76 (1988). These methods generally entail exposing a predetermined amount of a substrate to a predetermined amount of a enzyme preparation, crude

- 11 -

or purified. The rate of conversion of the substrate to the desired end product is measured under specified conditions of pH and temperature. Crude enzyme preparations can be produced by
5 culturing microorganisms in a suitable medium and then concentrating the enzyme that accumulates in the broth surrounding the cells using conventional means such as precipitation or ultrafiltration. The resulting enzyme preparations can be assayed for
10 specific activity by methods known to the art, including those that are keyed, respectively, to weight loss of insoluble substrates, changes in turbidity of polysaccharide suspensions, an increase in reducing end groups, a decrease in the viscosity
15 of polysaccharides such as beta-mannans, colorimetric determinations, measurements of clearance zones in polysaccharide-agar, or polarography.

In accordance with the present invention,
20 soil isolates are screened for hemicellulase production by culturing isolated colonies of the microorganisms in a hemicellulose-containing broth medium. After incubation, the broth medium is centrifuged and the resulting supernatant is
25 filtered to remove cells and to produce a crude preparation of hemicellulase. The enzyme is added to alkaline, viscous preparations of a preferred hemicellulose and the degree of liquefaction of the substrate over time is measured.

30 In a preferred embodiment of the present invention, soil isolates are cultured in a broth medium, wherein a mannan-containing hemicellulose is the sole carbon source. After incubation, the broth medium is centrifuged and the resulting supernatant
35 is filtered. The resulting crude enzyme preparation is added to a highly viscous preparation of a mannan-containing hemicellulose at a pH in the range

- 12 -

of 8-11. The relative activity of the enzyme is determined by the amount of time required to liquefy the hemicellulose.

5 In a particularly preferred embodiment of the present invention, soil isolates are cultured in a selective enrichment medium, wherein guar is the sole carbon source. After incubation, the culture is centrifuged and the resulting supernatant filtered to recover crude enzyme. The crude enzyme
10 is then introduced into tubes containing a cross-linked guar preparation, with metal ions added to enhance the viscosity of the preparation, at a pH in the range of 9-9.5. After incubation for at least one hour at 39-40° C, enzyme activity is measured by
15 determining the viscosity of a enzyme/guar solution using conventional viscosimetric procedures. See, e.g., *Biochim. Biophys. Acta* 139: 238 & 248 (1967); *Eur. J. Biochem.* 51: 207 (1975).

A hemicellulase obtained as described above
20 will generally display an activity characterized by a pH profile -- the curve defined by the relationship between a given enzymatic activity and pH -- that ranges over some 6.5 pH units, between minima at about pH 4.5 and about pH 11,
25 respectively. A pH profile of this sort is highly distinctive when compared, for example, to corresponding profiles for known *Bacillus mannanases*, which are virtually inactive at above about pH 8. See Araujo & Ward, *J. App. Bacteriol.*,
30 68: 253-61 (1990).

A hemicellulase used in the present invention preferably has a pH profile with a peak (i.e., the portion of the profile where activity is greatest) that is in a range from about pH 9 to above about
35 pH 7. It is also preferred that the enzyme display significant biological activity under conditions characterized both by high alkalinity and by

- 13 -

elevated temperatures. Such a suitable enzyme would be active, for example, when pH is in the range of 8-11 and the temperature is 60° C or higher.

5 A hemicellulase particularly preferred for use according to the present invention is an endo- β -D-mannanase having the following properties:

(1) Activity: A preferred enzyme acts on hemicellulosic materials containing mannan carbohydrates like galactomannan, glucomannan and
10 mannan. Activity in this context can be measured in the following way. An aqueous suspension containing 1.0% guar is used as a substrate, and 2 mL of 2 M glycine is added to 16.0 mL of the substrate. The mixture is mixed thoroughly and then preheated to 38°
15 C. Enzyme is added to the substrate, mixed well and then the viscosity of the enzyme/guar solution is measured using a simple approach, as by timing the charging and discharging of a pipette with stopwatch, or more complex devices like Brookfield
20 or Fann viscometers. A standard curve is prepared using a commercially available hemicellulase enzyme with known activity such as Miles B1500. Grams or units of the commercially available hemicellulase are plotted against the time required to change
25 viscosity and the new hemicellulase is compared to that of the commercial product.

(2) Substrate specificity: The enzyme degrades relatively simple carbohydrate polymers like guar gum, which is a polymer of D-mannose linked β -1,4
30 with D-galactose, and locust bean gum, as well as more complex, mannan-containing carbohydrates, for example, from soybean and alfalfa. Other suitable substrates will include mannan-containing coconut residue, carob bean gum, cassava, copra, and
35 chemically-modified forms of guar.

(3) Optimum pH: The optimum pH of the enzyme is about 7.0 or slightly above, e.g., in the range 7.1

- 14 -

to 7.5. The enzyme is stable in a pH range of about 4.5 to 11 (see Figure 4).

(4) Optimum temperature: The optimum temperature for enzyme activity is 40°C, but enzyme activity is observed at temperatures ranging from 20° to 90°C. As shown in Figure 2, the enzyme exhibits significant activity at temperatures ranging from 60° C to 90°C.

(5) Deactivation by Temperature and pH: At pH 9.0 the enzyme retains 50% of its maximum activity after 5.5 hours at 60° C, after 45 minutes at 75°C, and after 15 minutes at 90°C (see Figure 2).

(6) Molecular Weight: After purification to homogeneity, the enzyme has a molecular weight of about 32,000 as determined by SDS-polyacrylamide gel electrophoresis. To obtain a crude enzyme preparation from culture broth, cells and debris are first removed with a 0.1-micron filter. The enzyme is then concentrated with a membrane having a 10,000 molecular weight cut-off. Once the enzyme becomes quite concentrated, however, a good proportion of the enzyme is able to pass through the membrane. The enzyme in this final permeate can be concentrated using a membrane with a 5,000 molecular weight cut-off, or can be precipitated with three volumes of acetone. After the precipitate is centrifuged, the supernatant is siphoned off and the pellet resuspended in 50 mM phosphate buffer. This concentrate is dialyzed versus 10 mM phosphate buffer (pH 7.0) and then applied to a DEAE-Sephacel column. The enzyme is eluted with a solution gradient of sodium chloride of increasing ionic strength ranging from 0-1.25 M in 50 mM phosphate buffer (pH 7.0). Fractions are collected and tested for enzyme activity. The fractions exhibiting maximum activity are pooled and examined using SDS-polyacrylamide gel

- 15 -

electrophoresis and compared to protein molecular weight standards.

Pursuant to the present invention, commercially useful quantities of hemicellulase can be produced by culturing the foregoing hemicellulase-producing microorganisms using conventional fermentation techniques. In this context, "fermentation" is used broadly to refer to any controlled microbial action by which useful products are obtained from the substrate(s) of that action. In accordance with the present invention, fermentation can be carried out in a stirred-tank reactor. This type of reactor is essentially a closed cylindrical tank containing agitators, baffles, heat exchange coils, and automatic controls for temperature, air flow, pressure, pH and foaming. A fermenter of this sort would be charged with essential nutrients and a hemicellulase inducer. After sterilization, the reactor is inoculated with a culture consisting essentially of microorganisms that have been selected for their capacity to produce hemicellulase as described above.

Continuous culture can increase fermenter productivity by eliminating the downtime of batch cultures. But it is sometimes difficult to maintain sterility in large scale continuous cultures. Accordingly, batch fermentations which utilize the bacteria of the present invention are preferred.

In a preferred embodiment of this invention, a Gram-positive *Bacillus* strain like *B. lentis* CMG1240 is utilized to produce commercially useful quantities of the enzyme. A fermenter charged with a medium consisting of industrial-grade nutrients, glycerol (carbon source) and a mannan-containing hemicellulose, such as guar gum, is inoculated with a culture consisting essentially of a Gram-positive *Bacillus* strain as described above. The fermenter

- 16 -

is maintained at about 35° C (500-1,000 rpm agitation with an air flow between 0.4 and 1.6 vvm) for about 12 hours. The hemicellulose, such as guar, is added to the fermenter during the growth phase to further induce enzyme production. The fermentation is harvested after about 1 to 7.5 hours, and then enzyme activity is measured by viscometric means.

Enzyme activity of the fermentation product is assayed by centrifuging aliquots of the fermentation broth and testing the resultant supernatant. Aliquots of the supernatant (or dilutions thereof) are added to a guar solution and any reduction in the viscosity of the enzyme/guar solution is measured routinely by recording the time required for a constant bore pipette to discharge a predetermined volume, as described above in (1) of the preferred hemicellulase properties.

The enzyme is recovered from the fermentation broth by filtration through a 0.5-micron filter to remove cells followed by ultrafiltration using a 5,000 or 10,000 molecular weight cut-off ultrafiltration membrane. The concentrate is combined with three volumes of acetone at about 4° C to precipitate the enzyme. The precipitate is allowed to settle for approximately 24 hours before the supernatant is siphoned off. The precipitate is then centrifuged at about 4° C and the resultant pellet resuspended in phosphate buffer.

The preparation thus obtained represents a substantially pure form of the inventive enzyme, that is, a preparation in which virtually no other enzymatic activity is represented other than the hemicellulase activity described above. From such a preparation, other formulations can be made that consist essentially of the inventive enzyme; in other words, the enzyme may be present with other components of the formulation, including other

SUBSTITUTE SHEET

- 17 -

enzymes, that do not unduly hinder the desired hemicellulase activity. For example, a hemicellulase of the present invention may be included in a pulp-treating formulation which also
5 includes lignin-degrading enzymes.

In an alternative method for producing a hemicellulase within the present invention, DNA encoding hemicellulase could be isolated and used, via known procedures, to transform a suitable host
10 organism such that the enzyme is produced by the recombinant host in commercially useful amounts. Hemicellulase-encoding DNA could be isolated by screening nucleic acid libraries generated from microorganisms expressing a hemicellulase according
15 to the present invention. See, e.g., Ausubel et al., CURRENT PROTOCOLS IN MOLECULAR BIOLOGY at 5 & 6, John Wiley and Sons, New York (1987, 1990) (hereafter "Ausubel"). Such libraries would be screened by means of oligonucleotide probes that are
20 complementary to a polynucleotide encoding, for example, a portion of the N-terminus of a hemicellulase within the present invention. Illustrative of such an N-terminal portion is the amino acid sequence Ala-Ser-Gly-Phe-Tyr-Val-Xxx-Gly-
25 Thr-Ile-Leu-Xxx-Asp-Ser-Thr-Gly-Asn-Pro-Phe-Lys-Ile-Xxx-Gly-Xxx-Asn ["Xxx" denotes an indeterminant amino acid], as discussed in Example 12 below. See Ausubel at 6.

Alternatively, other portions including or
30 adjacent to the endogenous coding sequence of a hemicellulase according to the present invention can be used, when isolated using a probe as a template for generating other probes useful for isolating a hemicellulase-encoding polynucleotide according to
35 the present invention, e.g., based on the N-terminal sequence described above. Such a probe could be used in a known manner to screen a genomic or cDNA

- 18 -

library as described above, see Ausubel, *loc. cit.*, or to synthesize polymerase chain reaction (PCR) probes for use in amplifying a cDNA generated from an isolated RNA which codes for a hemicellulase of the present invention. Such a cDNA could then be cloned into a suitable expression vector and employed to transform a host organism. See Ausubel at §15.4.

A suitable polynucleotide in this regard would preferably comprise a nucleotide sequence, corresponding to the desired amino-acid sequence, that is optimized for the host of choice, as described below, with regard to codon usage, the initiation of translation, the most appropriate glycosylation pattern (if applicable), and the expression of recoverable amounts of a commercially useful hemicellulase within the present invention. Also, a vector selected for transforming the chosen host organism with such a polynucleotide molecule should allow for efficient maintenance and transcription of the sequence encoding the polypeptide. Such a vector is readily available or derivable from commercial sources, and is suited to a particular host cell employed for expressing a hemicellulase according to the present invention. For vectors exemplary of those suitably used to the present invention, see Ausubel at §§2-4.

Host cells suitable for expressing a hemicellulase according to the present invention would include prokaryotic or eukaryotic cells, for example, bacterial, algal, yeast, insect, mammalian and human cells. Thus, among host cells suitable for the present invention would be microorganisms of the genera *Aeromonas*, *Aspergillus*, *Bacillus*, *Echerichia*, *Kluyveromyces*, *Pichia*, *Rhodococcus*, *Saccharomyces* and *Streptomyces*. More specifically illustrative of the suitable host microorganisms

- 19 -

would be the bacteria *E. coli*, *B. subtilis*, *B. brevis* (J. Bacteriol. 172: 1312-20) and *B. lentis*, as well as the yeast *S. cerevisiae* strain X2181-1B, with genotype a *trp1 gall adel his2* (available from the Yeast Genetic Stock Center, Berkeley, California); strain ATCC 52683, with genotype a *his2 adel trp1 met14 ura 3* (available from the American Type Culture Collection, Rockville, Maryland); and strain ATCC 46183, with genotype a *his1 trp1* (also available from the American Type Culture Collection). Host cells comprising a polynucleotide encoding a hemicellulase according to the present invention can be grown under conditions that provide expression of a hemicellulase of the present invention in recoverable or commercially useful amounts. See, e.g., Ausubel at §§ 1 & 13.

A microorganism that produces a hemicellulase according to the present invention can be utilized to degrade a gelling agent used to fracture an oil well. More specifically, it is well known in the art to use mixtures of hydraulic gelling agents such as guar, proppants such as sand, glass bead and stones, and water in fracturing oil wells. Pumping such a mixture into a well at high pressure causes expansion of any natural fracture in a subterranean geologic formation adjacent to the well, resulting in a subsequent depositing of the mixture throughout the formation. But in order to obtain oil from the treated formation, the hydraulic gelling agent, preferably guar-based, must be removed.

Temperatures at the fracture site can range from ambient to 200° C and the ground water is generally alkaline with a pH range of 4.0 to 12.0 with a large percentage between pH 9.0 to 12.0. Pursuant to the present invention, a thermostable, alkaline-resistant hemicellulase can be utilized to degrade a guar-based hydraulic gelling agent

- 20 -

deposited in such a fracture. In a preferred embodiment, an effective amount of hemicellulase according to the present invention is added to a mixture of cross-linked guar and sand at the well head, immediately prior to the mixture being pumped down the well into the formation. Pressure is maintained on the well for 1-48 hours. In a controlled manner, the hemicellulase completely degrades the guar solution into small-molecule material of low viscosity, permitting extraction of oil from the newly fractured rock formation.

The present invention is further described below by reference to the following illustrative examples, in which the following media were used:

15 Selective Enrichment Broth (amount/liter)

10.0 g Guar Gum
5.0 g $(\text{NH}_4)_2\text{SO}_4$
pH 9.5

Selective Enrichment Agar (amount/liter)

20 2.0 g Guar Gum
 1.0 g Na_2HPO_4
 3.0 g $(\text{NH}_4)_2\text{SO}_4$
 0.2 g NaCl
 0.2 g $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$
25 50.0 mg $\text{CaCl}_2 \cdot 2\text{H}_2\text{O}$
 1.0 mL Trace Elements Solution I (see below)
 15.0 g Agar Noble
 50.0 mM Tris Buffer (pH 9.0)
 1.0 mL Vitamin Solution (see below)

30 Trace Elements Solution I (amount/liter)

100.0 mg EDTA
230.0 mg $\text{ZnSO}_4 \cdot 7\text{H}_2\text{O}$
180.0 mg $\text{MnSO}_4 \cdot \text{H}_2\text{O}$
60.0 mg H_3BO_3
35 100.0 mg $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$
 40.0 mg $\text{Na}_2\text{MoO}_4 \cdot 2\text{H}_2\text{O}$
 40.0 mg $\text{CoCl}_2 \cdot 6\text{H}_2\text{O}$
 70.0 mg KI
 40.0 mg $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$
40 0.4 mg $\text{NiCl}_2 \cdot 6\text{H}_2\text{O}$
 8.0 μL 0.1 M H_2SO_4

- 21 -

Vitamin Solution I (amount/liter)

5
1.0 g Vitamin B12
1.0 g Riboflavin, B2
1.0 g Pyridoxine, B6
1.0 g D-biotin
1.0 g Thiamine Hydrochloride
1.0 g Nicotinic Acid
1.0 g D-Ca-Pantothenic acid

Seed Broth (amount/liter)

10
7.5 g Glycerol
10.0 g Yeast Extract
2.5 g Corn Steep Liquor
1.0 g KH_2PO_4
2.0 g $(\text{NH}_4)_2\text{SO}_4$
15
0.5 g $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$
1.0 mL Trace Elements Solution II (see below)
pH 7.0-7.5

Fermentation Broth (amount/liter)

20
20.0 g Glycerol
20.0 g Yeast Extract
5.0 g Corn Steep Liquor
2.0 g KH_2PO_4
4.0 g $(\text{NH}_4)_2\text{SO}_4$
1.0 g $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$
25
2.0 mL Trace Elements Solution II
pH 8.5

Trace Elements Solution II (amount/liter)

30
20.0 g $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$
20.0 g $\text{FeCl}_3 \cdot 6\text{H}_2\text{O}$
0.5 g $\text{MnSO}_4 \cdot \text{H}_2\text{O}$
50.0 mg $\text{CoSO}_4 \cdot 7\text{H}_2\text{O}$
10.0 mg $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$
20.0 g $\text{CaCl}_2 \cdot \text{H}_2\text{O}$
50.0 mg H_3BO_3
35
100.0 g $\text{ZnSO}_4 \cdot 7\text{H}_2\text{O}$
100.0 g $\text{Na}_2\text{MoO}_4 \cdot 2\text{H}_2\text{O}$

Example 1: Isolation of hemicellulase-producing microorganisms from soil

40
Soil samples collected from both a tropical rain-forest and a temperate garden were added to selective enrichment broth in a 10% w/v

- 22 -

concentration. The cultures were shaken in baffled Erlenmeyer shake flasks at 37° C for 4 days. Four dilutions [1:10, 1:20, 1:800 and 1:50,000] of the initial cultures were made with fresh selective enrichment broth and incubated at 37° C for 4 days. The 1:50,000 dilution was used to prepare a dilution series (10^{-1} to 10^{-8}) in 0.85% NaCl which were plated out on selective enrichment agar and incubated at 34° C for 5-7 days. Following incubation, isolated colonies were selected from the plated cultures and streaked for purity three successive times on a suitable agar medium. Nine isolates from soil collected from a tropical rain-forest and 24 isolates from soil collected from a temperate garden were selected as a result of this screening process.

Example 2: Screening of soil isolates for hemicellulase activity

Each purified isolate from the tropical rain-forest soil was transferred to a shake flask containing selective enrichment broth and shaken for 48-72 hrs at 34° C. After incubation, the culture was centrifuged for 20-30 minutes at 10,000 rpm (4° C). The resultant supernatant was filtered sequentially through a 0.8- and a 0.45-micron filter to recover crude enzyme free of microbial cells. Each purified isolate from the temperate garden soil was transferred to tubes containing selective enrichment broth and shaken 12-48 hours at 34° C. After incubation, the cultures were centrifuged and filtered as described above. The crude enzyme solution was added to tubes containing 5 mL of a cross-linked guar preparation (5.0 g guar, 2.0 g $(\text{NH}_4)_2\text{SO}_4$ and 0.6 g sodium tetraborate per 400 mL water, pH 9.5) and incubated for at least 1 hour in a water bath at 39-40° C. To measure enzyme activity, a constant bore, 1-mL pipette was charged

- 23 -

with 1.0 mL of the mixture and allowed to discharge 0.9 mL. A stopwatch was used to measure the time required for this discharge to occur ("drop time"). A drop time of less than two seconds indicated measurable amounts of hemicellulase activity. Twelve of the twenty-four isolates recovered from temperate garden soil, and four of the nine isolates recovered from tropical rain-forest soil produced measurable amounts of hemicellulase.

10 Example 3: Production of commercially useful quantities of hemicellulase

A single, isolated colony of *B. lentis* (CMG1240) was inoculated into a 500 mL Erlenmeyer baffled shake (pre-seed) flask containing 50 mL of brain heart infusion broth. The inoculated flask was shaken (300 rpm) while incubating for 12-16 hours at 35° C. After incubation, the entire volume of the pre-seed flask was aseptically transferred to a 4 L Erlenmeyer baffled shake-flask (seed) containing one liter of seed medium. The seed-flask was incubated with shaking for 12-16 hours at 35° C. The entire volume of the seed flask was aseptically transferred to a 14-liter fermenter containing 9.0 liters of fermentation broth. The fermenter was maintained at 40° C, >20% dissolved oxygen and 500-1000 rpm. Guar gum was added to make a final concentration of 0.05% to the fermenter 2 or 3 times during the growth phase and once (0.5%) during the stationary phase to induce enzyme production. After 1-7.5 hours, the fermentation was terminated and enzyme activity assayed according to the procedure of Example 5. Assay data are shown in Figure 1.

- 24 -

Example 4: Measurement of hemicellulase activity to determine optimal culture conditions for hemicellulase production

Twenty-milliliter aliquots of 1% guar in water, mixed for 3 hours to obtain the maximum viscosity, and 2 mL aliquots of 2 M glycine-NaOH buffer (pH 9.0) were added to a glass-jacketed reactor maintained at 40° C by means of a circulating water bath. Aliquots of the fermentation product were centrifuged in a Microfuge for 5 minutes and the resultant supernatant was used to test for enzyme activity. A 100-200 mL sample of the supernatant was added to the reactor. Enzyme activity was measured as described in Example 5. Sample enzyme activity was calculated from a standard curve prepared by plotting the log of the drop time (seconds) of dilution of a standard hemicellulase (Miles B-1500) versus incubation time (minutes). The slope (on a linear scale) of the log plot was calculated and replotted versus grams or "chamber units" of hemicellulase.

Example 5: Assay for hemicellulase activity suitable for testing standards and purified enzyme preparations

In a procedure suitable for screening enzyme activity of standard hemicellulase or purified enzyme preparations, approximately 16.0 mL of a guar preparation, comprising 1% guar gum in water, and 2.0 mL of 2 M glycine (adjusted to pH 9.0 with NaOH), were added to a 100 mL glass jacketed reactor maintained at 38° C by means of a circulating water bath. The solution was mixed thoroughly by a magnetic stirrer after reaching temperature. Viscosity at time zero was determined by measuring the time required for the solution to drop from the 1.0 mL mark to the 0.7 mL mark in a constant bore, 1 mL pipette. The initial drop time of the guar

SUBSTITUTE SHEET

- 25 -

solution without enzyme added was 75 to 90 seconds. After enzyme addition (200-500 ML) to the reactor, the guar/enzyme solution was allowed to incubate. A time-zero measurement of the solution was immediately determined. Drop time measurements were performed until either 5 measurements had been made or until the drop time was less than 30 seconds. A standard curve was prepared by plotting the log of the drop time versus the time of incubation using different standard enzyme dilutions.

Example 6: Recovery of hemicellulase from fermentation

The procedure described in Example 3 was used in the production of useful quantities of hemicellulase in each of six 14-liter fermentations. After the fermentation was terminated, cells were removed from 60 L of culture broth by ultrafiltration with a 0.1 micron filter. Then, the enzyme was concentrated to a volume of approximately 3.5 L with a 10,000 molecular weight cut-off ultrafiltration membrane. The concentrate was combined with three volumes of acetone at 4° C to precipitate the hemicellulase. The precipitate was collected by allowing it to settle for 24 hours and then siphoning off the supernatant. The precipitate was then centrifuged at 6000 rpm for 10 minutes at 4° C. The resulting pellet was immediately resuspended in 170 mL of 50 mM potassium phosphate buffer (pH 7.0) to a final volume of 300 mL. The enzyme activity was determined according to the method of Example 5. The recovered hemicellulase had 9.45×10^6 chamber units/liter, which was equivalent to 6.3 kg/L of standard commercial hemicellulase (assayed at pH 9.0) or 1.89 kg/L total of standard hemicellulase formulation.

- 26 -

Example 7: Alternate process for recovering hemicellulase

Hemicellulase was recovered by the method described in Example 6, wherein methanol was used alternatively to precipitate hemicellulase. The supernatant was siphoned off and the precipitate centrifuged as described above. The paste-like precipitate was then blended with 10 volumes (w/w) of lactose and dried under vacuum at temperatures up to 40° C.

Example 8: Purification of hemicellulase

To produce a more purified product, two precipitation steps were used to obtain refined enzyme. A concentrate from ultrafiltration was prepared according to the method described in Example 6. The concentrate contained 6.39×10^6 units/liter, with a specific activity of 1.11×10^3 units/mg. Equal volumes (2.2 liters) of the concentrate and acetone were mixed to make a 50% (v/v) solution, and the resultant precipitate was then allowed to settle for 18 hours at 10° C. After the supernatant was collected, an additional volume of acetone was added to make the solution 75% (v/v). The resultant precipitate was allowed to settle for 24 hours at 10° C, after which the supernatant was first removed by siphoning and finally by centrifugation at 5000 rpm. The precipitate was then dissolved in 222 mL of 50 mM phosphate buffer (pH 7.0). Activity was determined to be 5.71×10^7 units/liter, with a specific activity of 6.17×10^3 units/mg. This purification step resulted in a recovery of 90% of the total activity with only 20% of the protein, causing a 5.56 fold increase in the specific activity.

SUBSTITUTE SHEET

- 27 -

Example 9: Alternative hemicellulase purification process

An alternative purification process was employed to avoid the precipitation step of Example 8. The hemicellulase was concentrated to the desired concentration by ultrafiltration as described in Example 6. A liquid formulation of the concentrate was then prepared by adding stabilizing and preservative agents such as ethylene glycol, or alternatively, glycerol, methanol, ethanol, sorbitol, sodium benzoate, sodium citrate, methylparaben, sodium ascorbate or an equivalent.

Example 10: Temperature stability of hemicellulase

An aliquot of broth was obtained from strain CMG1240 cultured in shake-flasks (48-72 hours at 35° C, pH 9.0) and clarified by centrifugation in a Microfuge. The supernatant was then filtered using a 0.45-micron filter; several aliquots were collected in sterile containers and stored on ice for use as controls. Some of these aliquots were subjected to steam sterilization for use as negative controls. The remaining supernatant was added to tubes and incubated in water baths maintained at 60° C and 75° C, and in a 90° C heating block. The activity of the enzyme at pH 9.0 and at these elevated temperatures over time was determined. The enzyme was stable at 90° C, exhibiting a half-life of 15 minutes in the absence of protecting agents. At 75° C, the enzyme reached 50% inactivation after 45 minutes. At 60° C, the enzyme reached 50% inactivation only after 5.5 hours. The results obtained in this experiment are presented in Figure 2.

Example 11: High temperature stability of hemicellulase formulated as a lactose powder

- 28 -

An aliquot of a hemicellulase solution prepared according to Example 8 (acetone precipitate) was extensively blended with 10 volumes (w/w) of powdered lactose. Two aliquots of the precipitate were dissolved in 10 mM phosphate buffer (pH 7.0) and stored on ice for use as controls. The dry lactose formulation was heated at 50° C for more than 400 hours. The test results, as shown in Figure 3, indicated that the lactose formulation showed virtually no loss of activity over the period tested.

Example 12: Characterization of the purified hemicellulase

A broth concentrate was prepared by ultrafiltration, i.e., cells were removed with a 0.1-micron membrane and the enzyme concentrated with a 10,000 molecular weight cut-off membrane. When the enzyme became concentrated, it was observed that some of the hemicellulase passed through the membrane.

This permeate fraction was re-concentrated with the 10,000 molecular weight cut-off membrane and then used for further purification. The re-concentrated permeate was precipitated by the addition of three volumes of acetone at 4° C. The supernatant was siphoned off, and the precipitate resuspended in 50 mM phosphate buffer (pH 7.0) to a final volume of 222 mL. The enzyme concentrate was subsequently dialyzed against 10 mM phosphate buffer (pH 7.0) to remove any salts. The dialyzed concentrate was then applied to a DEAE-Sephacel column (2.8 x 30 cm). Elution of the hemicellulase was carried out by adding a solution of increasing ionic strength to the column at a flow rate of 1 mL/minute. The solution consisted of 0-1.25 M NaCl in 50 mM phosphate buffer.

- 29 -

Ten-milliliter fractions were collected and tested for enzyme activity by measuring viscosity reduction of a 1% guar solution as described in Example 5. Fractions 96-109, which showed maximum activity, were pooled and then analyzed (along with molecular weight marker-proteins) on a 12.5% SDS-polyacrylamide gel. The results of this analysis indicated that the hemicellulase was highly purified, and had a molecular weight of 32,000. Hemicellulase fractions were ultimately purified by electrophoresis on the same type of SDS-gel and blotted onto Immobilon P PVDF membrane and stained with coomassie blue by the method of Matsudaira, *J. Biol. Chem.* 262: 10035-38 (1987). The hemicellulase band on the membrane was excised and used for amino acid sequence analysis. The enzyme was analyzed by Edman degradation using a Biosystems Gas Phase Sequenator. The N-terminal sequence of the purified enzyme was characterized as follows:

1	5	10
Ala-Ser-Gly-Phe-Tyr-Val-Xxx-Gly-Thr-Ile-Leu-Xxx-		
15	20	25
Asp-Ser-Thr-Gly-Asn-Pro-Phe-Lys-Ile-Xxx-Gly-Xxx-Asn		

Example 13: Percentage relative activity of hemicellulase vs. pH

A hemicellulase preparation was prepared by admixing 30.0 lb Hydroxypropyl Guar per 1,000 gallons of 2% KCl in water. A 20 mL portion of the preparation was placed in a Brookfield viscometer UL cup. A control was prepared by adding 1 mL of deionized water to the 20 mL guar preparation. The viscometer cup was then stoppered and shaken. The cup was placed on the viscometer and dial readings were measured at 3 rpm for 10 minutes. Approximately 6 minutes were required to achieve stable readings.

SUBSTITUTE SHEET

- 30 -

Twenty-milliliter aliquots of the gel preparation were adjusted to various pH levels.

One milliliter of broth from a shake flask as described in Example 2 was added to each aliquot.

5 The dial readings measured at 3 rpm after 10 minutes were subtracted from the control dial readings.

These differences were normalized relative to the results obtained at pH 7.0 at which maximum enzyme activity was observed. See Table II.

10 The plot of relative enzyme activity vs. pH demonstrates that the enzyme displayed greater than 90% activity over a pH range of 6 to 9. Figure 4 compares the activity of the enzyme vs that of a commercially available enzyme over a pH range of 2 to 11. The enzyme retained slightly more than 20% relative activity at pH 10.8.

TABLE II

% RELATIVE ACTIVITY OF HEMICELLULASE VS. PH

	<u>pH</u>	<u>Dial Reading at 3 rpm</u>	<u>Control-Sample Differential*</u>	<u>% Relative Activity</u>
20	10.80	42.5	9.0	21.4
	9.75	25.0	26.5	63.1
	9.17	17.5	34.0	92.9
	8.33	10.5	41.0	97.6
25	7.00	9.5	42.0	100.0
	6.25	11.0	40.5	96.4
	4.64	27.0	24.5	58.3

30 * Calculated by subtracting the dial reading of the sample from the dial reading (51.5) of the water control. Represents the loss in viscosity due to the presence of the enzyme.

Example 14: Use of hemicellulase under simulated field tests

35 A base gel was prepared by admixing the following in pounds per 1,000 gallons of 2% KCl in water:

- 31 -

40.0 lb Hydroxypropyl Guar
 2.5 lb Fumaric Acid
 10.0 lb Sodium Bicarbonate
 pH 7.4

- 5 The base gel corresponds to a commercial hydraulic gelling agent which would be used in the stimulation of a subterranean formation.

The base gel was admixed in a 4 liter Waring Blender for approximately 30 minutes after which
 10 time it was divided into 200 mL aliquots and permitted to further hydrate for at least 1 hour. Each sample was then further mixed with a variable speed laboratory stirrer equipped with a homogenizer blade for about 1 minute during which time the
 15 following were added to the fluid: 144 grams of sand (equivalent to 6 pounds per gallon concentration); enzyme formulated with lactose powder in a quantity identified below; and an organotitanate cross-linker in an amount sufficient
 20 to provide a concentration of 33 ppm Ti^{+6} . The base gel with additives was then placed in a heated bath maintained at 60° C. The viscosity of the cross-linked gel was measured with a Model 35 Fann Viscometer. The results, presented in Tables III
 25 and VI, demonstrate that a sufficient quantity of the hemicellulase will degrade cross-linked hydroxypropyl-guar fluid containing either sand or resin-coated sand.

TABLE III

30 20/40 Mesh U.S. Sieve Series, Ottawa Sand

Enzyme (lb/1000 gallons)

0 0.2 0.3 0.4 0.5 0.65 0.8 1.0

Time(hr)

Viscosity (centipoise)

1	C	C	C	11	7	-	-	4
35 2	C	C	WC	8	5	-	-	-
4	C	C	12	8	5	-	-	-
6	C	C	10	7	5	-	-	-
22	C	C	8	6	5	-	-	-

C= Cross-linked; WC= Weak Cross-link; - = No Data

SUBSTITUTE SHEET

- 32 -

TABLE VI

20/40 Mesh U.S. Sieve Series, Resin-Coated Sand

Enzyme (lb/1000 gallons)
0 0.2 0.3 0.4 0.5 0.65 0.8 1.0

5	<u>Time(hr)</u>	<u>Viscosity (centipoise)</u>							
		C	C	-	-	C	C	WC	24
	1	C	C	-	-	C	WC	30	20
	2	C	C	-	-	C	WC	20	10
	4	C	C	-	-	C	WC	10	7
	6	C	C	-	-	C	WC	7	6
10	22	C	C	-	-	C	WC		

C= Cross-linked; WC= Weak Cross-link; - = No Data

Example 15: Activity of the enzyme vs the pH of the base gel preparation.

A base gel was prepared by admixing the following in 1,000 gallons of 2% KCl in water:

40.0 lb Hydroxypropyl Guar

0.1 gallon ammonium/acetic acid buffer solution

10.0 lb Sodium Bicarbonate

The base gel was prepared according to the method described in Example 14 and maintained at 79.5° C in a heated water bath. The viscosity of the base gel preparation was measured according to the method described in Example 14. The results are presented in Table V and VI.

TABLE V

HEMICELLULASE VS. BASE GEL PH (8.0-8.5)

Dial Readings at 3 rpm

30	<u>Time(hr)</u>	<u>Control</u>	<u>0.25lb/1000 gal</u>	<u>0.50lb/1000 gal</u>
	0	35	32	32
	2.0	20	11	8
	6.0	19	10	8
	22.0	18	7	6
	Final pH	9.1	8.8	9.1

SUBSTITUTE SHEET

- 33 -

TABLE VI

HEMICELLULASE VS. BASE GEL PH (9.5-10.0)Dial Readings at 3 rpm

	<u>Time(hr)</u>	<u>Control</u>	<u>0.50lb/1000 gal</u>	<u>1.0lb/1000 gal</u>
5	0	34	34	34
	2.0	22	-	-
	3.0	-	18	16
	5.0	19	-	-
	7.0	-	17	15
10	22.0	18	18	12
	Final pH	10.1	9.5	9.5

Example 16: Characterization of *Bacillus lentis* CMG1240 by lipid analysis

An analysis was conducted of the fatty acid
 content of one of the microorganisms isolated
 according to the methods of Examples 1 and 2 above.
 Methods of fatty acid analysis used to characterize
 and identify microorganisms have been reported by E.
 Eerola and O.P. Lehtonen, *J. Clin. Microbiol.*
 26:1745-1753 (1988) and G.M. Mukwaya and D.F. Welch,
J. Clin. Microbiol. 27:2640-2646 (1989). These
 methods have been refined by Microbial ID, Inc.
 (Newark, Delaware) and Hewlett-Packard. The refined
 method, referred to as the MIDI microbial
 identification system, analyzes fatty acid methyl
 esters prepared from microbial lipids by gas-liquid
 chromatography. A microorganism isolated according
 to the methods of Examples 1 and 2 and designated
 strain CMG1240 was analyzed using the MIDI microbial
 identification system. The results of the analysis
 are presented in Table VII. In tests 1 through 4,
 the cells were grown at 28° C and pH 6.7 on TSBA agar
 according to the standard Microbial ID method. In
 test 5, the cells were grown at 35° C and pH 8.5,
 conditions which cause a significant change in lipid
 composition.

- 34 -

TABLE VII

PERCENTAGE OF TOTAL LIPIDS

<u>FATTY ACID NAME</u>	TEST 1	TEST 2	TEST 3	TEST 4	TEST 5
14:0 ISO	1.05	2.32	1.51	1.58	1.06
14:0	2.16	2.96	1.82	1.69	3.05
15:0 ISO	13.24	11.19	12.62	11.95	15.98
15:0 ANTEISO	42.06	42.41	42.15	42.95	35.42
15:0	1.16	1.56	0.95	-	2.42
16:0 ISO	2.28	4.42	3.65	4.30	3.42
16:1 A	9.62	6.88	6.98	6.28	4.35
16:0	17.69	21.11	16.68	18.91	20.82
17:1 ISO E	1.83	-	1.60	-	-
17:0 ISO	4.67	3.81	6.70	6.88	5.64
17:0 ANTEISO	3.79	3.33	5.32	5.45	3.96
18:1 C15 9	-	-	-	-	3.88

A comparison of the fatty acid profile presented in Table VII with profiles of all the strains of known *Bacillus* species in the MIDI fatty acid data base revealed that the subject profile most closely resembled profiles of microorganisms identified as *Bacillus lentis*.

SUBSTITUTE SHEET

- 35 -

WHAT IS CLAIMED IS

1. In substantially pure form, an enzyme that catalyzes the degradation of hemicellulose, wherein said enzyme has a pH profile for its activity in catalyzing said degradation that ranges from about pH 4.5 to about pH 11.
2. An enzyme according to Claim 1, wherein said profile peaks between about pH 7 and about pH 9.
3. An enzyme according to Claim 1, wherein catalysis by said enzyme occurs both (A) at a pH in the range of 8-11 and (B) at a temperature of at least 60° C.
4. An enzyme according to Claim 3, wherein said enzyme is a *Bacillus* enzyme.
5. An enzyme according to Claim 1, wherein said enzyme catalyzes the degradation of mannan-containing hemicellulose.
6. An enzyme according to Claim 5, wherein said enzyme is an endo- β -D-mannanase.
7. An enzyme preparation consisting essentially an enzyme that catalyzes the degradation of hemicellulose, wherein said enzyme has a pH profile for its activity in catalyzing said degradation that ranges from about pH 4.5 to about pH 11.

SUBSTITUTE SHEET

- 36 -

8. A microbial culture consisting essentially of bacterial cells that produce an enzyme that catalyzes the degradation of hemicellulose, wherein said enzyme has a pH profile for its activity in catalyzing said degradation that ranges from about pH 4.5 to about pH 11.

9. A microbial culture according to Claim 8, wherein said profile peaks between about pH 7 and about pH 9.

10. A microbial culture according to Claim 8, wherein catalysis by said enzyme occurs both (A) at a pH in the range of 8-11; and (B) at a temperature of at least 60° C.

11. A microbial culture according to Claim 8, wherein said bacteria are of the genus *Bacillus*.

12. A microbial culture according to Claim 11, wherein said bacteria are of the species *Bacillus lentis*.

13. A method of producing commercially useful quantities of hemicellulase using a microbial culture according to Claim 8, comprising the steps of

(a) growing said microbial culture in a suitable broth medium;

(b) adding a mannan-containing hemicellulose to said broth culture during the growth phase or the stationary phase of said microorganism;

(c) obtaining a culture supernatant from said broth culture, said supernatant comprising a

- 37 -

predetermined concentration range of said hemicellulase;

(d) removing said hemicellulase by ultrafiltration;

5 (e) precipitating said hemicellulase with a precipitating agent selected from the group consisting of acetone and methanol; and

(f) recovering said hemicellulase.

10 14. A method of conditioning subterranean oil wells comprising the steps of

(a) preparing a suspension comprised of water, a hydraulic gelling agent and a proppant, said proppant having an indicated mesh size;

15 (b) pumping an adequate amount of said suspension under high pressure into an oil well causing a fracture to occur in subterranean geologic formations adjacent to said well such that said suspension is deposited in said fracture; and

20 (c) pumping an adequate amount of an enzyme according to Claim 1 into said fracture with said suspension to effect degradation of said hydraulic gelling agent.

25 15. An isolated polynucleotide encoding an enzyme that catalyzes the degradation of hemicellulose, wherein said enzyme has a pH profile for its activity in catalyzing said degradation that ranges from about pH 4.5 to about pH 11.

30 16. A vector comprising a nucleotide sequence coding for an enzyme that catalyzes the degradation of hemicellulose, wherein said enzyme has a pH profile for its activity in catalyzing

- 38 -

said degradation that ranges from about pH 4.5 to about pH 11.

17. A host cell transformed with a vector according to Claim 16 such that said host cell
5 expresses said enzyme.

18. A method for producing an enzyme that catalyzes the degradation of hemicellulose, comprising the steps of

(A) providing a host cell according to
10 Claim 17,


(B) culturing said host cell under conditions wherein said host cell expresses said enzyme in recoverable amounts; and

(C) recovering said enzyme.

1 / 4

FIG. 1

14 L Production - Assay Data




- Fermentor #1
1/100 -200uL
- △ Fermentor #2
1/100 -200uL
- Fermentor #3
1/100 -200uL
- + Fermentor #4
1/100 -200uL
- Fermentor #5
1/100 -200uL

2 / 4

FIG. 2

Temperature Stability

pH 9.0


 Δ 90 C

○ 75 C


□ 60 C

8 / 4

FIG. 3



4 / 4



SUBSTITUTE SHEET

INTERNATIONAL SEARCH REPORT

International Application No. PCT/US91/03756

I. CLASSIFICATION OF SUBJECT MATTER (if several classification symbols apply, indicate all) *

According to International Patent Classification (IPC) or to both National Classification and IPC

IPC(5): C12N 9/00, 9/42

U.S. CL.: 166/244.1; 435/42, 183,209

II. FIELDS SEARCHED

Minimum Documentation Searched ⁷

Classification System	Classification Symbols
U.S. CL.:	435/42,183,209; 166/244.1

Documentation Searched other than Minimum Documentation
to the Extent that such Documents are Included in the Fields Searched *

III. DOCUMENTS CONSIDERED TO BE RELEVANT *

Category *	Citation of Document, ¹¹ with indication, where appropriate, of the relevant passages ¹²	Relevant to Claim No. ¹³
Y	Agric. Biol. Chem.. Volume 52, No. 3, issued 1988. T. Akino et al., "Characterization of Three -Mannanases of an Alkalophilic <u>Bacillus</u> sp.," pages 773-779, see entire document.	1-7
Y	Applied Microbiology and Biotechnology, Volume 26, issued 1987, T. Akino et al., "Production of -mannosidase and -mannanase by an alkalophilic <u>Bacillus</u> Sp.," pages 323-327, see entire document.	1-7,13
Y	US, A, 4,169,945 (DeGuia et al.) 02 October 1979, see column T, lines 15-24.	14
Y	US, A. 4,356,095 (Leavitt) 26 October 1982, see claims 1 and 2.	14

* Special categories of cited documents: ¹⁰

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step

"Y" document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

IV. CERTIFICATION

Date of the Actual Completion of the International Search

05 September 1991

Date of Mailing of this International Search Report

30 OCT 1991

International Searching Authority

ISA/US

Signature of Authorized Officer

Lori Yuan

ebw

This Page Blank (uspto)